

FIRST AND SECOND ORDER AHARONOV BOHM EFFECTS IN THE

EVANS UNIFIED FIELD THEORY.

by

Myron W. Evans,

Alpha Institute for Advanced Study (AIAS),

emyrone@aol.com

ABSTRACT

The first and second order Aharonov Bohm effects are explained straightforwardly in the Evans unified field theory using the spin connection generated by electromagnetism as spinning spacetime.

Keywords: Evans unified field theory; first and second order Aharonov Bohm effects.

1. INTRODUCTION.

The class of first order Aharonov Bohm (AB) effects {1} (those due to a static magnetic field) can be defined as AB effects in which the wavenumber (κ) of a matter beam such as an electron beam is shifted by the electromagnetic potential A acting at first order in the minimal prescription:

$$\kappa \rightarrow \kappa + \frac{e}{\hbar} A. \quad - (1)$$

Here $-e$ is the charge on the electron and \hbar the reduced Planck constant. Experiments on the AB effect can be summarized schematically with reference to Fig. (1), which defines one area within another as follows:



In the well known Chambers experiment {2} for example the outer area is that enclosed by the interacting electron beams in a Young diffraction set up, and the inner area is that enclosed by an iron whisker within which is trapped a static magnetic flux density B . In the standard model, electromagnetism always is a theory of special relativity and:

$$\underline{B} = \underline{\nabla} \times \underline{A} \quad - (2)$$

where \underline{A} is the vector potential. The AB effect is observed in the Chambers experiment as a shift in the diffraction pattern of the electron beams, a shift that is proportional to:

$$\underline{\Phi} = \int_S d \wedge A \quad - (3)$$

in which the surface integral is around the OUTER boundary defined by the paths of the two

electron beams. This is despite the fact that \underline{B} and therefore $\underline{\nabla} \times \underline{A}$ are confined to the INNER boundary $\{2, 3\}$ defined by the circumference of the iron whisker. The latter is placed between the openings of the Young interferometer. In the standard model, if B vanishes then so does $\frac{dA}{\lambda}$. This is clearly stated in a standard textbook such as ref. $\{2\}$. Therefore in the standard model there cannot be regions in which $\frac{dA}{\lambda}$ exists and in which B does not exist. Despite this simple inference it is often claimed confusingly that the first order AB effect is due to the effect of non-zero $\frac{dA}{\lambda}$ where B is zero or that the AB effect is a pure quantum effect with no classical counterpart. Other attempts $\{2\}$ at explaining the first order AB effect in the standard model rely on the classical concept of gauge transforming A. This confusion shows that the standard model does not explain the first order AB effect satisfactorily, or at all. This much is evidenced by over fifty years of theoretical controversy, all caused by the use of special relativity where general relativity is needed. The Evans field theory $\{3-6\}$ is the first successful unified field theory that develops electromagnetism unified with gravitation as a correctly objective field theory of general relativity.

In Section 2 it is argued that the gauge transform theory of the standard model violates Stokes' Theorem in non-simply connected regions, and so is erroneous and unable to explain correctly the first order AB effect. In Section 3, the first order AB effect is explained correctly and straightforwardly using the spin connection of the Evans field theory. The latter is therefore preferred experimentally and mathematically to the standard model. Finally in Section 4 the second order or electromagnetic Aharonov Bohm effect is explained through the conjugate product of potentials in the Evans field theory, a conjugate product that defines the well known Evans spin field and which is observed in the inverse Faraday effect IFE $\{7\}$. The IFE is explained from the first principles of general relativity in the Evans unified field theory $\{3-6\}$ but cannot be explained in the standard model without the empirical or ad hoc introduction of the conjugate product $\{8\}$ in non-linear optics. Similarly for the

second order AB effect which is implied by the well observed IFE.

2. ARGUMENT AGAINST THE STANDARD MODEL

Adopting the well known { 9 } notation of differential geometry the

following three equations summarize the attempted description of the first order Aharonov

Bohm effect in the standard model:

$$F = d \wedge A \quad - (4)$$

$$d \wedge F = 0 \quad - (5)$$

$$k \rightarrow k + \frac{e}{\hbar} A. \quad - (6)$$

Experimentally the observed Aharonov Bohm effect in an experiment such as that of

Chambers is proportional to the magnetic flux (in weber) within the outer boundary of Fig (1)

(the boundary defined by the paths of the electron beams):

$$\underline{\Phi} = \int_S d \wedge A = \int_S F = \oint A \text{ (outer boundary)}. \quad - (7)$$

However, the magnetic flux density of the iron whisker is at the same time confined within

the inner boundary

$$\underline{\Phi} = \int_S d \wedge A = \int_S F = \oint A \text{ (inner boundary)} \quad - (8)$$

and \underline{A} is also confined within the inner boundary in the standard model. There is a contradiction between Eqs. (7) and (8) because the experimentally measured flux is given by Eq. (7) but the physical magnetic flux is given by Eq. (8). In a standard model textbook such as ref. { 2 }, pp. 101 ff. an attempt is made to explain this contradiction in the first order Aharonov Bohm effect using the gauge transformation:

$$A \rightarrow A + d\chi. \quad - (9)$$

The standard model uses the Stokes Theorem to argue that:

$$\oint d\chi \neq 0 \quad (?) \quad - (10)$$

in the region between the inner and outer boundary of Fig (1) and that the Aharonov Bohm effect is due to the integral over $d\chi$ in Eq. (10). However, the basis of electromagnetic gauge theory in the standard model is the Poincaré Lemma:

$$d \wedge (d\chi) := 0 \quad - (11)$$

which is true for simply AND multiply connected spaces. The integrated form of the Poincaré Lemma is the Stokes Theorem:

$$\underline{\Phi} = \int_S F = \int_S d \wedge (d\chi) = \oint d\chi := 0 \quad - (12)$$

which is also true for multiply connected spaces { 10 }. The standard model { 2 } attempts to explain the first order AB effect by asserting INCORRECTLY that:

$$\underline{\Phi} = \int_S F = \int_S d \wedge (d\chi) = \oint d\chi \neq 0. \quad (?) \quad - (13)$$

In order to apply the Stokes Theorem to Fig (1) for example, a cut { 10 } is made to join the outer and inner boundaries as follows:




Fig. (2)

and contour integration proceeds in one direction around the inner boundary, across the cut, in the opposite direction around the outer boundary, and back across the cut. Examples of such procedures are to be found in a standard textbook on vector algebra { 10 }, in problems on the application of the Stokes Theorem.

We must look to general relativity and the Evans unified field theory for first correct explanation of the first order Aharonov Bohm^m effect.

3. EXPLANATION OF THE FIRST ORDER AB EFFECT IN EVANS THEORY.

In the correctly objective description of the first order AB effect { 11 } the electromagnetic field is defined by the first Maurer Cartan structure equation:

$$F^a = D \wedge A^a = d \wedge A^a + \omega^a_b \wedge A^b$$

- (14)

where D^{\wedge} is the covariant exterior derivative, d^{\wedge} is the exterior derivative, ω^a_b is the spin connection in the well known Palatini formulation of general relativity { 12, 13 } in which the tetrad e^a_{μ} is the fundamental field. (In the original Einstein Hilbert formulation of general relativity the metric is the fundamental field.) The electromagnetic potential field is the fundamental tetrad field within a primordial or universal scalar $A^{(0)}$, where $cA^{(0)}$ has the units of volts, and where c is the speed of light in vacuo:

$$A_{\mu}^a = A^{(0)} q_{\mu}^a. \quad - (15)$$

It is seen that this gives a natural field unification scheme, because the metric used by Einstein and Hilbert is well known { 9, 14 } to be the dot product of two tetrads:

$$g_{\mu\nu} = q_{\mu}^a q_{\nu}^b \eta_{ab} \quad - (16)$$

where η_{ab} is the Minkowski metric of the tangent bundle whose index is a. The latter becomes essentially a polarization index { 3 - 6 } in the Evans field theory. For example:

$$a = (1), (2), (3) \quad - (17)$$

describes circular polarization where ((1), (2), (3)) is the well known { 15 } complex circular basis. Again, it is well known { 16 } that the tetrad is developed into the spin 3 / 2 gravitino in supersymmetry theory, and that the Einstein Hilbert and Palatini variations of general relativity are inter-related by the tetrad postulate { 9, 16 }:

$$D_{\nu} q_{\mu}^a = 0. \quad - (18)$$

One of the major inferences of the Evans field theory is that the tetrad field is the fundamental entity of objective (i.e. generally covariant) unified field theory, a unified field theory which satisfies the fundamental requirements of objectivity and general covariance in physics, the principles of general relativity. Electromagnetism in the standard model is a theory of special relativity, and is Lorentz covariant only. So the standard model is not a correctly objective theory of physics. This is the fundamental reason why it cannot describe the first order Aharonov Bohm effect, and gauge theory in special relativity { 2 } suffers from the same fundamental defect.

From Eq. (14) the magnetic flux in weber from the Evans field theory is defined

as:

$$\overline{\Phi}^a = \int_S F^a = \oint A^a + \int_S \omega^a{}_b \wedge A^b \quad - (19)$$

and is in general the sum of two terms, one involving the spin connection $\omega^a{}_b$ of general relativity. It is $\omega^a{}_b$ that gives rise to the first (and second) order Aharonov Bohm effects. The fundamental reason is that the second term on the right hand side of Eq. (19) exists in the outer region of Fig (1) even though the magnetic flux density F^a is confined to the inner region and so is zero in the region between the inner and outer boundaries. The second term on the right hand side of Eq. (19) does not vanish, and gives rise to the AB effects. In the Chambers experiment, for example, the observed shift in the electron diffraction pattern is:

$$\delta = \alpha \int_S \omega^a{}_b \wedge A^b \quad - (20)$$

where α is a proportionality constant. The integration in Eq. (20) is around the outer boundary as required experimentally, the boundary defined by the diffracting electron beams in the Young interferometer of the Chambers experiment. The latter therefore observes the spin connection of the Evans theory directly. The spin connection is not present in the standard model, which has no explanation (Section 2) for the AB effects. The spin connection is a direct consequence of the major discovery of the Evans field theory that electromagnetism is the spinning of spacetime {3-6} ¹⁷ - the spinning spacetime gives rise directly to the spin connection in the Palatini variation of general relativity.