

THE NEW GENERAL RELATIVITY: SELF CONSISTENCY AND FIELD
EQUATION

by

M. W. Evans and H. Eckardt,
Civil List, AIAS and UPITEC.

(www.webarchive.org.uk, www.aias.us, www.atomicprecision.com, www.upitec.org,
www.et3m.net)

ABSTRACT

Using a cyclic permutation of three chain rule equations the new general relativity is shown to be rigorously correct and self consistent. The orbital Evans identities are computed and shown to be two simultaneous equations which describe the time evolution of the orbit. These reduce to a simple differential equation which can be solved by computer.

Keywords: ECE theory, the new general relativity, the Evans orbital identities, time evolution and field equation.

4FT 207

1. INTRODUCTION

In recent papers of this series {1 - 10} it has been shown that the Einsteinian general relativity is trivially incorrect in many ways, and it has been replaced by efforts towards forging a new relativity based on special relativity developed straightforwardly into general relativity in the context of orbital dynamics. The method employed is to constrain the Minkowski metric with any observed orbit, so the new relativity is perfectly general. As shown in Section 2 of this paper the constraint reduces the dimensionality of the metric, and introduces a mathematical space in which Riemann torsion and curvature are non-zero. In the unconstrained space of the Minkowski metric of special relativity, all elements of torsion and curvature are zero. The infinitesimal line element may be deduced rigorously from the constrained metric. The constraint is the observed orbit itself, so in the new relativity the orbit is analysed in terms of torsion and curvature. There is one independent torsion element and two independent curvature elements for each observed orbit. So an ephemeris of all cosmology may be built up using torsion and curvature. In section 3 the field equation and time evolution equation of any orbit is deduced from the Evans identity {1 - 10}, which is a fundamental identity of geometry deduced during the evolution {1 - 10} of the ECE theory.

2. METRIC, LINE ELEMENT AND CONNECTIONS

Consider the Minkowski line element of special relativity in the plane:

$$d\tau^2 = 0 \quad \text{--- (1)}$$

of any planar orbit. In cylindrical polar coordinates {11} the line element ds^2 is related to the metric $g_{\mu\nu}$ by:

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = c^2 dt^2 - dr^2 - r^2 d\theta^2 \quad - (2)$$

where

$$g_{00} = 1, \quad g_{11} = -1, \quad g_{22} = -1, \quad - (3)$$

$$dx^0 = c dt, \quad dx^1 = dr, \quad dx^2 = r d\theta.$$

Here c is the vacuum speed of light, assumed to be a constant. The cylindrical polar coordinates in the plane (1) are denoted (r, θ) , and t is time. Any planar orbit is described by the orbital equation:

$$\dot{r} = \frac{dr}{dt} = \dot{r}(r(t), \theta(t)) \quad - (4)$$

in which $r(t)$ and $\theta(t)$ are both functions of time. Therefore the infinitesimal line element of any planar orbit may be written as:

$$ds^2 = c^2 dt^2 - dr^2 - \left(r \frac{d\theta}{dr}\right)^2 dr^2 \quad - (5)$$

in which:

$$dx^2 = \left(r \frac{d\theta}{dr}\right) dx^1 \quad - (6)$$

and:

$$ds^2 = g_{00} dx^0 dx^0 + g_{11} dx^1 dx^1 + g_{22} \left(r \frac{d\theta}{dr}\right)^2 dx^1 dx^1. \quad - (7)$$

Note carefully that the number of coordinates have been reduced from three to

two. The spacetime becomes non-Minkowski in the sense that:

$$ds^2 = g_{00} dx^0 dx^0 + g_{11} dx^1 dx^1 \quad - (8)$$

$$g'_{11} = g_{11} + f g_{22} \quad - (9)$$

where the function f is defined by:

$$f = \left(r \frac{d\theta}{dr} \right)^2 \quad - (10)$$

The metric of the new general relativity is therefore:

$$g_{\mu\nu} = \begin{bmatrix} 1 & 0 \\ 0 & -(1+f) \end{bmatrix} \quad - (11)$$

Previous work {1 - 10} has shown that the connection is antisymmetric, and that

the metric compatibility equation defines the one independent connection of any orbit as:

$$\Gamma^1_{01} = \frac{1}{2c g_{11}} \frac{\partial g_{11}}{\partial t} \quad - (12)$$

with, by definition of antisymmetry:

$$\Gamma^1_{01} = -\Gamma^1_{10} \quad - (13)$$

In order to evaluate the connection consider the function f written out as follows in three cyclic permutations. If:

$$f = f(r(t), \theta(t)) \quad - (14)$$

then by the chain rule {12} of differentiation:

$$\frac{df}{dt} = \frac{\partial f}{\partial r} \frac{dr}{dt} + \frac{\partial f}{\partial \theta} \frac{d\theta}{dt} \quad - (15)$$

Similarly, if:

$$f = f(t(\theta), r(\theta)) \quad - (16)$$

then:

$$\frac{df}{d\theta} = \frac{\partial f}{\partial r} \frac{dr}{d\theta} + \frac{\partial f}{\partial t} \frac{dt}{d\theta} \quad - (17)$$

Finally if:

$$f = f(\theta(r), t(r)) \quad - (18)$$

then:

$$\frac{df}{dr} = \frac{\partial f}{\partial \theta} \frac{d\theta}{dr} + \frac{\partial f}{\partial t} \frac{dt}{dr} \quad - (19)$$

From Eqs. (15) and (19):

$$\frac{df}{dt} = \frac{\partial f}{\partial r} \frac{dr}{dt} + \frac{d\theta}{dt} \frac{dr}{d\theta} \left(\frac{df}{dr} - \frac{\partial f}{\partial t} \frac{dt}{dr} \right) \quad - (20)$$

so:

$$\frac{df}{dt} = \frac{\partial f}{\partial r} \frac{dr}{dt} \quad - (21)$$

From Eqs. (15) and (17) it may be shown similarly that:

$$\frac{df}{d\theta} = \frac{\partial f}{\partial \theta} \frac{d\theta}{dt} \quad - (22)$$

From Eqs. (17) and (21):

$$\frac{\partial f}{\partial t} = \frac{d\theta}{dr} \frac{dr}{dt} \left(\frac{df}{d\theta} - \frac{\partial f}{\partial t} \frac{dt}{d\theta} \right) = \frac{df}{dt} - \frac{\partial f}{\partial t} \quad (23)$$

therefore:

$$\frac{\partial f}{\partial t} = \frac{1}{2} \frac{df}{dt} \quad (24)$$

From Eqs. (15) and (22):

$$\frac{\partial f}{\partial t} = \frac{d\theta}{dt} \frac{dr}{d\theta} \left(\frac{df}{dr} - \frac{\partial f}{\partial t} \frac{dt}{dr} \right) = \frac{df}{dt} - \frac{\partial f}{\partial t} \quad (25)$$

so the same result as Eq. (24) is obtained self consistently:

$$\frac{\partial f}{\partial t} = \frac{1}{2} \frac{df}{dt} \quad (26)$$

Finally use:

$$\frac{df}{dt} = \frac{df}{d\theta} \frac{d\theta}{dt} \quad (27)$$

to find that:

$$\frac{\partial f}{\partial t} = \frac{1}{2} \omega \frac{df}{d\theta} \quad (28)$$

The function

$df/d\theta$

can be found from any observed orbit, and the angular velocity:

$$\omega = \frac{d\theta}{dt} \quad (29)$$

can be observed by astronomy. So the connection (12) can be found for any orbit.

3. TORSION, CURVATURE AND EVANS IDENTITY.

As in previous work and using computer algebra any orbit in the plane (1) is due to the connection:

$$\Gamma^1_{01} = -\Gamma^1_{10} = \frac{df/dt}{2c(1+f)} \quad - (30)$$

The single independent element of torsion of any planar orbit is twice the connection:

$$T^1_{01} = -T^1_{10} = \frac{df/dt}{c(1+f)} \quad - (31)$$

For any planar orbit of the type (1) there are two independent elements of curvature:

$$R^1_{001} = -R^1_{010} = -\frac{2 \left((1+f) d^2f/dt^2 - (df/dt)^2 \right)}{2c^2(1+f)^2} \quad - (32)$$

and:

$$R^1_{101} = -R^1_{110} = \frac{(df/dt)(df/dr) - (1+f) d^2f/dr dt}{2c(1+f)^2} \quad - (33)$$

Finally there are two equations of the Evans identity {1 - 10}:

$$D_0 T^1_{01} = R^1_{001} \quad - (34)$$

and

$$D_1 T^1_{10} = R^1_{110} \quad - (35)$$

for any planar orbit of cosmology. Computer algebra evaluates these to be:

$$6(1+f) \frac{d}{dt} \left(\frac{df}{dt} \right) = 5 \left(\frac{df}{dt} \right)^2 \quad - (36)$$

and

$$(1+f) \frac{d}{dr} \left(\frac{df}{dt} \right) = \frac{df}{dr} \frac{df}{dt} \quad - (37)$$

These must be solved simultaneously. Denoting:

$$x = \frac{df}{dt}, \quad y = \frac{df}{dr} \quad - (38)$$

then dividing Eq. (36) by Eq. (37) it is found that:

$$6 \frac{dx}{dt} = 5 \frac{dx}{dt} \quad - (39)$$

Therefore:

$$\frac{d}{dt} \left(\frac{df}{dt} \right) = 0 \quad - (40)$$

This is a type of conservation law for any planar orbit. It is a time evolution equation:

$$\frac{d}{dt} \left(\omega \frac{df}{d\theta} \right) = \frac{d\omega}{dt} \frac{df}{d\theta} + \omega \frac{d}{dt} \left(\frac{df}{d\theta} \right) = 0 \quad - (41)$$

which is true for any planar orbit. Its solution is:

$$\omega \frac{df}{d\theta} = A \text{ (no explicit } t \text{ dependence)} \quad - (42)$$

an equation which may be integrated to give f in terms of two constants A and B :

$$f = \int \frac{A}{\omega} d\theta. \quad - (43)$$

The second constant B is a constant of integration.

ACKNOWLEDGMENTS

The British Government is thanked for a Civil List Pension and the staff of AIAS and others for many interesting discussions. Dave Burleigh is thanked for posting, Alex Hill, Robert Cheshire and Simon Clifford for translation and broadcasting. The AIAS is established under the aegis of the Newlands Family Trust (2011).

REFERENCES

- {1} M. W. Evans, Ed., J. Found. Phys. Chem., (CISP, Cambridge, www.cisp-publishing.com) from June 2011 in six issues a year.
- {2} M. W. Evans, "Definitive Refutations of the Einsteinian General Relativity", Issue Six of reference (1), softback and e book, (CISP, Spring 2012, ISBN 978-1-907343-08-7 (softback) and ISBN 9781-907343-52-0 (e book)).
- {3} M. W. Evans, S. Crothers, H. Eckardt and K. Pendergast, "Criticisms of the Einstein Field Equation" (CISP, Spring 2011).
- {4} M. W. Evans, H. Eckardt and D. W. Lindstrom, "Generally Covariant Unified Field Theory" (Abramis Academic, Suffolk, 2005 - 2011) in seven volumes.
- {5} L. Felker, "The Evans Equations of Unified Field Theory" (Abramis 2007, translated into Spanish by Alex Hill on www.aias.us, open source).
- {6} K. Pendergast, "The Life of Myron Evans" (CISP, 2011).
- {7} The ECE websites, www.webarchive.org.uk, www.aias.us, www.upitec.org.

www.et3m.net, www.atomicprecision.com, archived in the National Library of Wales, British Library, and the US, Russian and European web archives supported by Library of Congress, N.S.F. and other organizations in San Francisco, Paris and Amsterdam.

{8} M .W. Evans and J. - P. Vigiér, "The Enigmatic Photon" (Kluwer, 1994 to 2002, hardback and softback), in ten volumes.

{9} M .W. Evans and S. Kielich, Eds., "Modern Nonlinear Optics" (Wiley, 1992, reprinted 1993, softback 1997, second edition and e book 2001), in six volumes and two editions,

{10} M .W. Evans and L. B. Crowell, "Classical and Quantum Electrodynamics and the B(3) Field" (World Scientific 2001).

{11} E. J. Milewski (chief editor), "Vector Analysis Problem Solver" (Staff of the Research and Education Association, New York, 1987).

{12} G. Wilkinson, "Mathematical Methods for Science Students" (Longmans, London, 1968).